

DYNAMICAL STABILITY OF FLUID SPHERES IN SPACETIMES WITH A NONZERO COSMOLOGICAL CONSTANT*

STANISLAV HLEDÍK[†], ZDENĚK STUHLÍK[‡] and KRISTINA MRÁZOVÁ[⊙]

*Institute of Physics, Faculty of Philosophy and Science, Silesian University in Opava,
Bezručovo nám. 13, Opava, CZ-746 01, Czech Republic*

E-mail: [†]Stanislav.Hledik@fpf.slu.cz, [‡]Zdenek.Stuchlik@fpf.slu.cz, [⊙]Kristina.Mrazova@fpf.slu.cz

The Sturm–Liouville eigenvalue equation for eigenmodes of the radial oscillations is determined for spherically symmetric perfect fluid configurations in spacetimes with a nonzero cosmological constant and applied in the cases of configurations with uniform distribution of energy density and polytropic spheres. It is shown that a repulsive cosmological constant rises the critical adiabatic index and decreases the critical radius under which the dynamical instability occurs.

Keywords: Perfect fluid configurations; Cosmological constant; Dynamical stability.

1. Introduction

The internal Schwarzschild spacetimes with nonzero cosmological constant ($\Lambda \neq 0$) and uniform distribution of energy density were given for star-like configurations¹ and extended to more general situations.² The polytropic and adiabatic spheres were preliminary treated and compared,^{3,4} neutron star models with regions of nuclear matter described by different relativistic equations of state that are matched were also treated.⁵ Their stability can be considered on energetic grounds⁶ or it can be treated in dynamical way.⁷ Here we determine the dynamical stability conditions for the uniform density and polytropic spheres using the approach of Misner et al.⁸

2. Sturm–Liouville Equation

Interior of a spherically symmetric configuration is described (in standard Schwarzschild coordinates) by the line element

$$ds^2 = -e^{2\Phi} dt^2 + e^{2\Psi} dr^2 + r^2(d\theta^2 + \sin^2\theta d\varphi^2), \quad (1)$$

with metric coefficients taken in the general form

$$\Psi = \Psi(r, t), \quad \Phi = \Phi(r, t). \quad (2)$$

The perfect fluid distribution is given by energy density and pressure profiles $\rho(r, t)$ and $p(r, t)$. The static equilibrium configuration is given by $\Phi_0(r)$, $\Psi_0(r)$, $\rho_0(r)$, $p_0(r)$ satisfying the Euler equations. The radially pulsating configuration is then determined by

$$q(r, t) = q_0(r) + \delta q(r, t), \quad \delta q \equiv (\delta\Phi, \delta\Psi, \delta\rho, \delta p, \delta n), \quad (3)$$

*This research has been supported by Czech grant MSM 4781305903.

where n is the number density. The pulsation is given by the radial displacement $\xi = \xi(r, t)$. The Euler perturbations δq are related to the Lagrangian perturbations measured by an observer who moves with the pulsating fluid by the relation

$$\Delta q(r, t) = q(r + \xi(r, t), t) - q_0(r) \approx \delta q + q'_0 \xi. \quad (4)$$

Introducing a new variable $\zeta \equiv r^2 e^{-\Phi_0} \xi$, the radial pulsations are governed by

$$W\ddot{\zeta} = (P\zeta')' + Q\zeta \quad (5)$$

with the functions $W(r)$, $P(r)$, $Q(r)$ determined for the equilibrium configuration

$$W \equiv (\rho_0 + p_0) \frac{1}{r^2} e^{3\Psi_0 + \Phi_0}, \quad (6)$$

$$P \equiv \gamma p_0 \frac{1}{r^2} e^{\Psi_0 + 3\Phi_0}, \quad (7)$$

$$Q \equiv e^{\Psi_0 + 3\Phi_0} \left[\frac{(p'_0)^2}{\rho_0 + p_0} \frac{1}{r^2} - \frac{4p'_0}{r^3} - (\rho_0 + p_0) \left(\frac{8\pi G}{c^4} p_0 - \Lambda \right) \frac{e^{2\Psi_0}}{r^2} \right]. \quad (8)$$

The linear stability analysis can be realized by the standard assumption of the displacement decomposition

$$\zeta(r, t) = \zeta(r) e^{i\omega t}. \quad (9)$$

Then the dynamic equation reduces to the Sturm–Liouville equation and the related boundary conditions in the form

$$(P\zeta')' + (Q + \omega^2 W)\zeta = 0, \quad (10)$$

$$\frac{\zeta}{r^3} \text{ finite as } r \rightarrow 0, \quad \gamma p_0 \frac{e^{\Phi_0}}{r^2} \zeta' \rightarrow 0 \text{ as } r \rightarrow R. \quad (11)$$

The Sturm–Liouville equation (10) and the boundary conditions determine eigenfrequencies ω_j and corresponding eigenmodes $\zeta_i(r)$, where $i = 1, 2, \dots, n$. The eigenvalue Sturm–Liouville (SL) problem can be expressed in the variational form⁸ as the extremal values of

$$\omega^2 = \frac{\int_0^R (P\zeta'^2 - Q\zeta^2) dr}{\int_0^R W\zeta^2 dr} \quad (12)$$

determine the eigenfrequencies ω_i and the corresponding functions $\zeta_i(r)$ are the eigenfunctions that have to satisfy the orthogonality relation

$$\int_0^R e^{3\Psi_0 - \Phi_0} (p_0 + \rho_0) r^2 \xi^{(i)} \xi^{(j)} dr = 0. \quad (13)$$

3. Results and Conclusions

We applied the Sturm–Liouville approach to spheres with uniform energy density,¹ and polytropic spheres.³ The case of uniform spheres can be properly taken as a test bed of the dynamical instability problem — although these solutions of the Einstein equations are of rather artificial character, they reflect quite well the basic properties




Figure 1. Dependence of the critical value of adiabatic index γ_c on sphere radius R . *Full curve*: vanishing cosmological constant $\lambda = 0$; then γ_c diverges as $R \rightarrow 9r_g/8$ from above. *Dashed curve*: positive cosmological constant $\lambda = 0.1$, the point of divergence is shifted to $1.18421 > 9/8$, and $\gamma_{c,\lambda>0} > \gamma_{c,\lambda=0}$. *Dashed-dotted curve*: negative cosmological constant $\lambda = -0.1$, the point of divergence is shifted to $1.07143 < 9/8$, and $\gamma_{c,\lambda<0} < \gamma_{c,\lambda=0}$.

of very compact objects.⁹ The dependence of the critical value of adiabatic index $\gamma \equiv (\partial \ln p / \partial \ln n)_s = (n/p)(\Delta p / \Delta n)$ on configuration radius R is for the uniform case given in Fig. 1 (see also¹²). The polytropic case is treated in details by Stuchlík and Hledík.¹⁰

Bibliography

1. Z. Stuchlík, *Acta Phys. Slovaca* **50**, 219 (2000).
2. C. G. Böhmmer, *Gen. Relativity Gravitation* **36**, p. 1039 (2004).
3. Z. Stuchlík and S. Hledík, General Relativistic Polytropes with a Nonzero Cosmological Constant, In preparation, (2006).
4. S. Hledík, Z. Stuchlík and K. Mrázová, General relativistic adiabatic fluid spheres with a repulsive cosmological constant, In preparation, (2006).
5. M. Urbanec, J. Miller and J. R. Stone, Matching of equations of state: influence on calculated neutron star models pp. 357–362. In Hledík and Stuchlík.¹¹
6. R. F. Tooper, *Astrophys. J.* **140**, 434 (1964).
7. S. Chandrasekhar, *Astrophys. J.* **140**, p. 417 (1964).
8. C. W. Misner, K. S. Thorne and J. A. Wheeler, *Gravitation* (Freeman, San Francisco, 1973).
9. N. K. Glendenning, *Phys. Rev. C* **37**, p. 2733 (1988).
10. Z. Stuchlík and S. Hledík, Radial pulsations and dynamical stability of spherically symmetric perfect fluid configurations in spacetimes with a nonzero cosmological constant pp. 209–222. In Hledík and Stuchlík.¹¹
11. S. Hledík and Z. Stuchlík (eds.), *Proceedings of RAGtime 6/7: Workshops on black holes and neutron stars, Opava, 16–18/18–20 September 2004/2005* (Silesian University in Opava, Opava, 2005).
12. C. G. Böhmmer and T. Harko, Dynamical instability of fluid spheres in the presence of a cosmological constant, *Phys. Rev. D* **71**, p. 084026 (2005), arXiv: gr-qc/0504075.